

Relaxed Gabor expansion at critical density and a ‘certainty principle’

V.P.Palamodov
Tel Aviv University

31.07.05

1 Introduction

The functions in $\mathbb{R}^n$ that are the best localized in configuration and in frequency spaces simultaneously are known in quantum field theory as coherent states (shifted modulated gaussian functions). Any coherent state gives exact minimum for the uncertainty relation which means that each coherent state $\mathbf{e}_\lambda$ is maximally concentrated at a certain point λ of the phase space $\Phi = \mathbb{R}^n \times (\mathbb{R}^n)'$. Another advantage of the family $\{\mathbf{e}_\lambda, \lambda \in \Phi\}$ is its large symmetry group: the Weyl-Heisenberg group is acting by shifts and modulation operators and the metaplectic group which is a ‘quantization’ of the group of linear symplectic transformations.

The role of the Weyl-Heisenberg family in the information theory ($n = 1$) was emphasized by Gabor [6]: “each elementary signal conveys exactly one datum, or one quantum of information”. Gabor’s idea was to expand an ‘arbitrary’ signal in a series of coherent states $\mathbf{e}_\lambda$ for points λ in a maximally sparse lattice Λ in the phase plane. The property of the Gabor system $\{\mathbf{e}_\lambda, \lambda \in \Lambda\}$ depends on the area a of a cell of the lattice Λ . The system is complete, if and only if $a \leq 1$. Lyubarskii [9] has shown that the Gabor system is a frame, if $a < 1$. We focus on the critical case $a = 1$ when the Gabor system of critical density is still complete, but is not a frame. A function $f \in L_2(\mathbb{R})$ may not have a convergent Gabor series, but there exists always a series that converges in a distribution sense, see Janssen [7]. The book of I.Daubechies [2] contains a survey of the problem, see also Feichtinger, Strohmer [4] for applications of the Gabor analysis.

Lyubarskii and K.Seip [10] has shown that for an arbitrary function $f \in L_2$ that is $1/2$ -smooth in time and frequency there exists a convergent series over a Gabor system, if a minor deformation of a lattice Λ of critical density is made.

We show here that any δ -smooth (in Φ) function for $\delta > 1$ has a *unique* convergent Gabor expansion with l_2 -coefficients, provided one more point $\sharp$ is added to

a lattice Λ ; this point is in the middle of a cell and is called *sharp* point. We call it a *relaxed* Gabor expansion. This approach can save Gabor's idea. The rate of convergence of the coefficients can be improved, if few more sharp terms are added to the Gabor system. The coefficients of such a series are again uniquely defined for a sufficiently smooth function f .

The relaxed Gabor expansion sheds light to the physical wisdom: *a field supported in certain domain D in the phase space has about $|D|$ degrees of freedom*, where $|D|$ means the symplectic area of the domain (Nyquist, Wigner, Brillouin, Shannon, Gabor,...). This claim can be called 'certainty' principle as opposite of the classical uncertainty one. The Landau-Pollak dimension theorem [8] gives an accurate form to this principle in a special situation. Given positive numbers T, Ω , any function f 'concentrated' in the box D of the size $T \times 2\Omega$ can be approximated by a linear combination of $2T\Omega + O(\log T\Omega)$ first prolate spheroidal functions. The condition of concentration means that the tail of a unit energy function f is small in $\mathbb{R} \setminus [0, T]$ and the tail of Fourier transform $\hat{f}$ is small in $\mathbb{R}^* \setminus [-\Omega, \Omega]$. The number $2T\Omega$ is equal to the area $|D|$ of the box measured by the canonical symplectic form in the phase plane. This means the approximation 'dimension' of the space of functions localized in the box equals the symplectic area of this up to a logarithmic term.

We formulate and prove here a rigorous form of the certainty principle for domains of arbitrary shape. We use Gabor means $\langle f | \mathbf{e}_\mu \rangle$ to specify the condition of concentration of a function $f \in L_2$ in the phase space Φ . We say that f is concentrated in a domain D , if the integral over $\Phi \setminus D$ of the density $|\langle f | \mathbf{e}_\mu \rangle|^2 d\mu$ is small. We show that for an arbitrary sets $K \subset D$, where D is r -neighborhood of K for some $r \geq r_0$, any function f concentrated in D can be written in the form $g + h + \phi$, where g is a linear combination of Gabor functions $\mathbf{e}_\lambda$, with $\lambda \in \Lambda \cap D$, h belongs to the linear envelope of Gabor functions with sharp $\lambda \in D \setminus K$ and the norm of ϕ is bounded by the rate of concentration of f plus an exponentially small term as $r \rightarrow \infty$. The total number N of such points λ is equal to $|D| + |D \setminus K| = |D| + O(r|D|^{1/2})$. To summarize briefly, this means that any function f concentrated in D is a linear combination of $N \sim |D| + O(r|D|^{1/2})$ coherent states supported in D up to a exponentially small term.

I thank Yu.Lyubarskii for helpful discussions.

2 Gabor transform and localization

Fix a coordinate x in a line $\mathbb{R}$; the family of functions in $\mathbb{R}$

$$\mathbf{e}_\lambda(x) \doteq 2^{1/4} \exp(-\pi(x-p)^2 + 2\pi i \theta x)$$

are called Gabor functions. Here $\lambda = (p, \theta)$ is a point in the phase space $\Phi \doteq \mathbb{R} \times \mathbb{R}'$ ($\mathbb{R}'$ means the dual line). Another term for $\mathbf{e}_\lambda$ is *coherent state* at the point λ . The phase space has the natural Euclidean structure: $|\lambda|^2 = |p|^2 + |\theta|^2$ and Lebesgue measure $d\lambda \doteq dp d\theta$. We use the notation $\langle \cdot | \cdot \rangle$ for the scalar product in $L_2 = L_2(\mathbb{R})$ and $\|\cdot\|$ for the norm. For $f \in L_2$ the function $\lambda \mapsto \langle f | \mathbf{e}_\lambda \rangle$ defined in Φ is called *Gabor transform* of f . This transform is unitary:

Proposition 2.1 *For an arbitrary function $f \in L_2$ the equation holds*

$$\int_{\Phi} |\langle f | \mathbf{e}_\lambda \rangle|^2 d\lambda = \|f\|^2 \quad (1)$$

where $d\lambda = dp d\theta$ and

$$f = \int_{\Phi} \langle f | \mathbf{e}_\lambda \rangle \mathbf{e}_\lambda d\lambda \quad (2)$$

in weak L_2 -sense.

PROOF. By Plancherel Theorem

$$\begin{aligned} \int_{\mathbb{R}} |\langle f | \mathbf{e}_\lambda \rangle|^2 d\theta &= 2^{1/2} \int \left| \int_{\mathbb{R}} f(x) \exp(-\pi(x-p)^2) \exp(2\pi i \theta x) dx \right|^2 d\theta \\ &= 2^{1/2} \int |F(f_p)(-\theta)|^2 d\theta = 2^{1/2} \int |f_p(x)|^2 dx, \end{aligned} \quad (3)$$

where $f_p(x) = f(x) \exp(-\pi(x-p)^2)$. Next we integrate both sides against dp and changing the variable x to $y = x - p$. This yields

$$\begin{aligned} \int \int |f_p(x)|^2 dx dp &= \int \int |f(x) \exp(-\pi y^2)|^2 dx dy \\ &= \int |f(x)|^2 dx \int \exp(-2\pi y^2) dy \\ &= 2^{-1/2} \int |f(x)|^2 dx \end{aligned}$$

and (1) follows. Now calculate the scalar product of integral (2) with $\mathbf{e}_\mu$. We have for arbitrary points $\lambda = (p, \theta), \mu = (q, \eta) \in \Phi$

$$\langle \mathbf{e}_\lambda | \mathbf{e}_\mu \rangle = \exp(\pi i (p+q)(\theta - \eta) - \pi |\lambda - \mu|^2 / 2). \quad (4)$$

Therefore

$$\int_{\Phi} \langle f | \mathbf{e}_\lambda \rangle \langle \mathbf{e}_\lambda | \mathbf{e}_\mu \rangle d\lambda = \langle f | E_{\lambda, \mu} \rangle,$$

where

$$\begin{aligned}
\bar{E}_{\lambda,\mu} &\doteq \int_{\Phi} \langle \cdot | \mathbf{e}_{\lambda} \rangle \langle \mathbf{e}_{\lambda} | \mathbf{e}_{\mu} \rangle d\lambda \\
&= \int_{\Phi} \langle \cdot | \mathbf{e}_{\lambda} \rangle \exp \left(\pi i (p+q) (\theta - \eta) - \pi |\lambda - \mu|^2 / 2 \right) d\lambda \\
&= 2^{1/4} \int \int \exp \left(-\pi (x-p)^2 - 2\pi i \theta x \right) \exp \left(\pi i (p+q) (\theta - \eta) - \pi |\lambda - \mu|^2 / 2 \right) d\theta dp
\end{aligned}$$

Calculate the interior integral by changing θ to $\xi = \theta - \eta$:

$$\begin{aligned}
\bar{E}_{\lambda,\mu}(x) &= 2^{1/2} \exp(-2\pi i \eta x) \int \exp \left(-\pi (x-p)^2 - \frac{\pi}{2} (p+q-2x)^2 - \frac{\pi}{2} |p-q|^2 \right) dp \\
&= 2^{1/2} \exp(-2\pi i \eta x) \exp(-\pi (x-q)^2) \int \exp(-2\pi (x-p)^2) dp = \bar{\mathbf{e}}_{\mu}(x)
\end{aligned}$$

This allows to write

$$\int_{\Phi} \langle f | \mathbf{e}_{\lambda} \rangle \langle \mathbf{e}_{\lambda} | \mathbf{e}_{\mu} \rangle d\lambda = \langle f | \mathbf{e}_{\mu} \rangle$$

This yields for an arbitrary $g \in L_2$

$$\langle f | g \rangle = \lim_{r \rightarrow \infty} \int_{|\lambda| \leq r} \langle f | \mathbf{e}_{\lambda} \rangle \langle \mathbf{e}_{\lambda} | g \rangle d\lambda = \int_{\Phi} \langle f | \mathbf{e}_{\lambda} \rangle \langle \mathbf{e}_{\lambda} | g \rangle d\lambda$$

The integral converges, since both factors belong to $L_2(\Phi)$. ►

Now we show that a function $f \in L_2$ can be localized in a convex domain $D \subset \Phi$ in terms of its Gabor transform.

Corollary 2.2 *For an arbitrary $f \in L_2$ and any $q \in \mathbb{R}$ we have*

$$\int_{p \geq q} |\langle f | \mathbf{e}_{\lambda} \rangle|^2 d\lambda = \int_{-\infty}^{\infty} I(x-q) |f(x)|^2 dx$$

where

$$I(x) \doteq 2^{1/2} \int_{-\infty}^x \exp(-2\pi y^2) dy$$

Remark. The function $I(x)$ tends fast to 1 and 0 as $x \rightarrow \infty$, respectively, $x \rightarrow -\infty$ and $I(0) = 1/2$.

PROOF follows from (3). ►

Corollary 2.3 *Let D be a domain and S be a rotation in the phase plane Φ and M_S be the corresponding metaplectic transform. We have*

$$\int |M_S f(x)|^2 I(x-q) dx \leq \int_{\Phi \setminus D} |\langle f | \mathbf{e}_{\lambda} \rangle|^2 d\lambda \tag{5}$$

where $q = \sup \{p; \lambda = (p, \theta) \in S(D)\}$.

PROOF. By Proposition 2.2 we have

$$\int |M_S f(x)|^2 I(x-q) dx \leq \int_{\Phi \setminus S(D)} |\langle M_S f | \mathbf{e}_\lambda \rangle|^2 d\lambda.$$

Because of the transform M_S is unitary, we have by Corollary 11.2 we have

$$\langle f | \mathbf{e}_\lambda \rangle = \langle M_S f | M_S \mathbf{e}_\lambda \rangle = \exp(-i\phi) \langle M_S f | \mathbf{e}_{S(\lambda)} \rangle$$

for some real phase ϕ . Therefore

$$\int_{\Phi \setminus S(D)} |\langle M_S f | \mathbf{e}_\lambda \rangle|^2 d\lambda = \int_{\Phi \setminus D} |\langle M_S f | \mathbf{e}_{S(\lambda)} \rangle|^2 d\lambda = \int_{\Phi \setminus D} |\langle f | \mathbf{e}_\lambda \rangle|^2 d\lambda$$

which completes the proof. ►

Suppose that the right-hand side of (5) is small. Taking in account that $I(x) \geq 1/2$ for $x \geq 0$, we conclude that for any rotation S the function $M_S f$ is strongly concentrated in the interval $[r, q] \doteq \pi(S(D))$, where $\pi : \Phi \rightarrow \mathbb{R}$ is the orthogonal projection. The inverse is true if, for instance, D is a convex polygon.

3 Gabor series

Choose some elements $\varepsilon \in \mathbb{R}$, $\varepsilon^* \in \mathbb{R}'$ such that $\varepsilon^*(\varepsilon) = 1$ and consider the lattice $\Lambda \subset \Phi = \mathbb{R} \times \mathbb{R}'$ generated by ε and ε^* :

$$\Lambda \doteq \{\lambda = k\varepsilon + j\varepsilon^*, k, j \in \mathbb{Z}\}.$$

The symplectic area of a cell is equal to $\varepsilon^*(\varepsilon) = 1$. Introduce a coordinate x in $\mathbb{R}$ such that $x(\varepsilon) = 1$; then we have $\xi(\varepsilon^*) = 1$ for the dual coordinate ξ in $\mathbb{R}'$. Consider the Gabor system $\{\mathbf{e}_\lambda, \lambda \in \Lambda\}$.

Proposition 3.1 *For an arbitrary numerical sequence $\{c_\lambda\} \in l_2(\Lambda)$ the series*

$$f = \sum_{\lambda} c_\lambda \mathbf{e}_\lambda$$

converges in L_2 and the inequality holds

$$\|f\| \leq \sigma_0 \left(\sum |c_\lambda|^2 \right)^{1/2}, \quad \sigma_0 = \sum \exp\left(-\frac{\pi}{2} k^2\right).$$

PROOF. By (4) we have for any $\lambda, \mu \in \Lambda$, $\lambda = (p, \theta)$

$$\begin{aligned} \langle \mathbf{e}_\lambda | \mathbf{e}_\mu \rangle &= g_{\lambda-\mu}, \\ g_\lambda &\doteq \exp\left(\pi i p \theta - \frac{\pi}{2} (p^2 + \theta^2)\right). \end{aligned} \tag{6}$$

Therefore

$$\|f\|^2 = \sum_{\lambda, \mu} c_\lambda \bar{c}_\mu \langle \mathbf{e}_\lambda | \mathbf{e}_\mu \rangle = \sum_{\lambda} c_\lambda \sum_{\mu} \bar{c}_\mu g_{\lambda-\mu}$$

The interior sum is a convolution and we can estimate its l_2 -norm as follows:

$$\sum_{\lambda} \left| \sum_{\mu} \bar{c}_\mu g_{\lambda-\mu} \right|^2 \leq \left(\sum |g_\lambda| \right)^2 \sum |c_\mu|^2 = \sigma_0^4 \sum |c_\lambda|^2,$$

since $\sum |g_\lambda| = \sigma_0^2$. This yields

$$\|f\|^4 \leq \sum_{\lambda} |c_\lambda|^2 \sum_{\lambda} \left| \sum_{\mu} \bar{c}_\mu g_{\lambda-\mu} \right|^2 \leq \sigma_0^4 \left(\sum |c_\mu|^2 \right)^2. \quad \blacktriangleright \quad (7)$$

4 Zak transform

The Zak transform of a function $f \in L_2$ is defined by the series

$$Zf(y, \xi) = \sum_{q \in \mathbb{Z}} \exp(2\pi i q \xi) f(y + q),$$

which converges almost everywhere in the square $\mathbb{Q} \doteq \{0 \leq y, \xi \leq 1\}$. The function Zf fulfils

$$Zf(y, \xi + 1) = Zf(y, \xi), \quad Zf(y + 1, \xi) = \exp(-2\pi i \xi) Zf(y, \xi). \quad (8)$$

Proposition 4.1 *The Zak transform is a unitary operator $L_2(\mathbb{R}) \rightarrow L_2(\mathbb{Q})$.*

PROOF is straightforward and we omit it. The inversion formula reads

$$\begin{aligned} f(x) &= \sum_{r \in \mathbb{Z}} \int_0^1 \int_0^1 dy d\xi g(y, \xi) \exp(2\pi i (y - x)(\xi + r)) \\ &= \int_0^1 dy \int_{\mathbb{R}} g(y, \xi) \exp(2\pi i \xi (y - x)) d\xi, \end{aligned}$$

where $g = Zf$ is 1-periodic function of ξ .

The Weyl-Heisenberg group acts in L_2 by the shifts and modulation operators

$$T_\lambda f(x) = \exp(2\pi i \theta x) f(x - p), \quad \lambda = (p, \theta) \quad (9)$$

We have for any $\lambda \in \Lambda$ we have

$$Z(T_\lambda f)(y, \xi) = \exp(2\pi i (p\xi + \theta\xi)) Zf(y, \xi). \quad (10)$$

The sum

$$\Theta(z) = 2^{1/4} \sum_{q \in \mathbb{Z}} \exp(2\pi i q z - \pi q^2)$$

is a Jacobi elliptic function: $\Theta(z) = 2^{1/2} \theta_3(z; \iota)$. This function is holomorphic of z in the whole plane, satisfies the periodicity conditions

$$\Theta(z+1) = \Theta(z), \quad \Theta(z+\iota) = \exp(\pi - 2\pi i z) \Theta(z) \quad (11)$$

has simple zero at the point $1/2 + \iota/2$ and no other zeros in the closed square $\mathbb{Q}$. The Zak transform of Gabor function for $\lambda = (p, \theta) \in \Lambda$ is expressed in terms of Θ :

$$Z\mathbf{e}_\lambda(y, \xi) = \exp(2\pi i(p\xi + \theta y)) \exp(-\pi y^2) \Theta(\xi + \iota y). \quad (12)$$

5 Creation and annihilation operators

The operator

$$\mathbf{a} = \frac{1}{2\pi} \frac{d}{dx} + x, \quad \mathbf{a}^+ = -\frac{1}{2\pi} \frac{d}{dx} + x$$

in L_2 are adjoint one to another. They are called the *annihilation* and the *creation* operators. Any Gabor function is an eigenvector of the annihilation operator:

$$\mathbf{a}\mathbf{e}_\lambda = \lambda \mathbf{e}_\lambda \quad (13)$$

where $\lambda = (p, \theta)$ and $\lambda = p + \iota\theta$. For any ϕ in the domain of the operator $\mathbf{a}$ we have

$$Z(\mathbf{a}\phi) = AZ\phi, \quad A \doteq \frac{1}{2\pi i} \left(\frac{\partial}{\partial \xi} + \iota \frac{\partial}{\partial y} \right) + y. \quad (14)$$

Consider the line bundle $\mathbf{E}$ over the torus $\mathbb{R}^2/\Lambda$ whose sections s fulfil the periodicity conditions

$$s(y, \xi + 1) = s(y, \xi), \quad s(y + 1, \xi) = \exp(-2\pi i \xi) s(y, \xi). \quad (15)$$

in the unit square $\mathbb{Q}$ (which is the fundamental domain of the lattice Λ). Let $W_2^\alpha(\mathbf{E})$ be the Sobolev space of order $\alpha \geq 0$ of sections ϕ of $\mathbf{E}$. By (8) the function Zf is a section of $\mathbf{E}$ of the class $W_2^0(\mathbf{E}) = L_2(\mathbb{Q})$. The differential operator A acts on smooth sections of $\mathbf{E}$ and defines for any α a bounded map $A : W^{\alpha+1}(\mathbf{E}) \rightarrow W^\alpha(\mathbf{E})$.

Proposition 5.1 *For any function $f \in L_2$ such that*

$$\int \left(|\lambda|^\delta + 1 \right) |\langle f | \mathbf{e}_\lambda \rangle|^2 d\lambda < \infty, \quad \delta > 0 \quad (16)$$

we have $Zf \in W_2^\delta(\mathbf{E})$.

PROOF. By (2) we can write

$$Zf = \int_{\Phi} \langle f | \mathbf{e}_{\lambda} \rangle Z\mathbf{e}_{\lambda} d\lambda$$

We have for arbitrary $\lambda = (p, \theta) \in \Phi$

$$Z\mathbf{e}_{\lambda}(\xi, y) = 2^{1/4} \sum_{q \in \mathbb{Z}} \exp(2\pi i q \xi + \theta(y + q)) \exp(-\pi(y + q - p)^2)$$

Setting $p = k + r, \theta = j + \eta, k, l \in \mathbb{Z}$ and $q' = q - k$ yields

$$\begin{aligned} Z\mathbf{e}_{\lambda}(y, \xi) &= 2^{1/4} \sum_{q \in \mathbb{Z}} \exp(2\pi i [q\xi + \theta y + q\eta]) \exp(-\pi(y + q - k - r)^2) \\ &= 2^{1/4} \exp(2\pi i [k(\xi + \eta) + \theta y]) \sum_{q \in \mathbb{Z}} \exp(2\pi i q'(\xi + \eta)) \exp(-\pi(y + q' - r)^2) \\ &= \exp(2\pi i [k\xi + jy]) \exp(2\pi i k\eta) Z\mathbf{e}_{(r, \eta)}(y, \xi) \end{aligned} \quad (17)$$

We can write

$$Zf(y, \xi) = \sum_{\mathbb{Z} \times \mathbb{Z}} \exp(2\pi i [k\xi + jy]) \int_{\mathbb{Q}} \langle f | \mathbf{e}_{\lambda} \rangle E_{\lambda}(y, \xi) dr d\eta,$$

The kernel $E_{\lambda}(y, \xi) \doteq \exp(2\pi i k\eta) Z\mathbf{e}_{(r, \eta)}(y, \xi)$ is a real analytic function for $\lambda, (y, \xi) \in \mathbb{R} \times \mathbb{R}$, is 1-periodic in ξ and its derivatives in y, ξ are bounded in any strip $|y| \leq C$ uniformly for $\lambda \in \Phi$. Take a test function ϕ in $\mathbb{R} \times \mathbb{R}$ and apply the Fourier transform $F = F_{(y, \xi) \mapsto (s, t)}$:

$$F(\phi Zf)(s, t) = \int_{\mathbb{Q}} \sum_{k, j} \langle f | \mathbf{e}_{\lambda} \rangle F(\phi E_{\lambda})(s - k, t - j) dr d\eta$$

The interior sum can be written as convolution on $\mathbb{Z}^2$:

$$G_{r, \eta} \doteq \sum_{k, j \in \mathbb{Z}} \langle f | \mathbf{e}_{(k, j) + (r, \eta)} \rangle F(\phi E_{\lambda})(s - k, t - j)$$

We have

$$|F(\phi E_{\lambda})(s, t)| \leq C_n (|s| + |t| + 1)^{-n}$$

for any natural n , since $\phi E_{\lambda} \in C^{\infty}$. Therefore for any fixed $(r, \eta) \in \mathbb{Q}$

$$\int (|s| + |t| + 1)^{\delta} |G_{r, \eta}(s, t)|^2 ds dt \leq C \sum_{k, j} (|k| + |j| + 1)^{\delta} |\langle f | \mathbf{e}_{(k, j) + (r, \eta)} \rangle|^2.$$

where the sum in the right-hand side converges for almost all (r, η) and the constant C does not depend on f . Integrating over $\mathbb{Q}$ yields

$$\begin{aligned}
\int (|s| + |t| + 1)^\delta |F(\phi Z f)(s, t)|^2 ds dt &= \int (|s| + |t| + 1)^\delta \left| \int_{\mathbb{Q}} G_{r, \eta}(s, t) dr d\eta \right|^2 ds dt \\
&\leq \int_{\mathbb{Q}} \int (|s| + |t| + 1)^\delta |G_{r, \eta}(s, t)|^2 ds dt dr d\eta \\
&\leq C \int_{\mathbb{Q}} \sum_{k, j} (|k| + |j| + 1)^\delta |\langle f | \mathbf{e}_{(k, j) + (r, \eta)} \rangle|^2 dr d\eta \\
&= C \int_{\Phi} (|k| + |j| + 1)^\delta |\langle f | \mathbf{e}_\lambda \rangle|^2 dp d\theta \\
&\leq C' \int_{\Phi} (|\lambda| + 1)^\delta |\langle f | \mathbf{e}_\lambda \rangle|^2 d\lambda < \infty
\end{aligned}$$

This implies that $\phi Z f$ belongs to W_2^δ and our statement follows. ►

Definition. Denote by H^δ the space of functions $f \in L_2$ that satisfies the condition (16) and set

$$\|f\|_\delta = \left(\int (|\lambda|^\delta + 1) |\langle f | \mathbf{e}_\lambda \rangle|^2 d\lambda \right)^{1/2}$$

Denote $H \doteq \cup_{\delta > 1} H^\delta$.

Remark 1. The space H^δ coincides with the modulation spaces $\mathbf{M}_{2,2}^w$ of H. Feichtinger, [3] for the weight function $w(\lambda) = |\lambda|^\delta + 1$.

Remark 2. The domain of the harmonic oscillator operator

$$H \doteq \frac{1}{2} (\mathbf{a}^+ \mathbf{a} + \mathbf{a} \mathbf{a}^+) = -\frac{1}{4\pi^2} \frac{d^2}{dx^2} + x^2.$$

is equal to the space H^2 .

6 Relaxed Gabor expansion

We show now that an arbitrary function $f \in H$ can be expanded in a Gabor series as above with one more term.

Denote $\sharp = (1/2, 1/2) \in \Phi$; we call *sharp* point any element $\mu \in \Lambda + \sharp$,¹. The

¹The motivation of this term comes from the interpretation of the lattice Λ as a table for a regular time-frequency notation of music (Brillouin-Wigner). The point $\sharp$ is then a tune which is shifted inside a rhythm interval (syncope), whose height is shifted by half a tune (sharp= "diese" tune).

sharp Poisson functional is the series

$$\gamma^\sharp(f) \doteq \frac{1}{i\Theta(0)} \sum_{\mathbb{Z}} (-1)^q f(q + 1/2) = \frac{Zf(\sharp)}{i\Theta(0)}.$$

We show below that this functional is well defined and continuous in H . In particular, we have $\gamma^\sharp(\mathbf{e}_\sharp) = 1$, whereas $\gamma^\sharp(\mathbf{e}_\lambda) = 0$ for arbitrary $\lambda \in \Lambda$, since Θ vanishes at each sharp point. We call the set $\Lambda^\sharp \doteq \Lambda \cup \{\sharp\}$ the *relaxed* lattice and consider the relaxed Gabor system $\{\mathbf{e}_\lambda, \lambda \in \Lambda^\sharp\}$.

Theorem 6.1 *There exists a family of continuous functionals $\gamma^\lambda, \lambda \in \Lambda$ in the space H such that an arbitrary $f \in H$ is developed in the series*

$$f = \sum_{\Lambda^\sharp} \gamma^\lambda(f) \mathbf{e}_\lambda \quad (18)$$

that converges in $L_2(\mathbb{R})$ and for any $\delta > 1$ there exists a constant C_δ such that

$$\sum_{\Lambda^\sharp} |\gamma^\lambda(f)|^2 \leq C_\delta \|f\|_\delta^2. \quad (19)$$

Lemma 6.2 *For an arbitrary $f \in H$ the Zak transform Zf is ε -Hölder continuous for $\varepsilon < \delta - 1$.*

PROOF OF LEMMA. By Proposition 5.1 $Zf \in W_2^\delta(\mathbf{E})$ for some $\delta > 1$. Sobolev's imbedding theorem implies that Zf is a continuous section of $\mathbf{E}$ and $\sup |Zf| \leq C \|f\|_\delta$, [11]. Moreover, Zf belongs to the Hölder ε -class for $\varepsilon < \delta - 1$. ►

PROOF OF THEOREM. By Lemma 6.2 the functional $\gamma^\sharp$ is well defined in H . Set $f_\sharp = f - \gamma^\sharp(f) \mathbf{e}_\sharp \in H$ and define $F \doteq Zf_\sharp / Z\mathbf{e}_0$. By Lemma 6.3 the function F is square integrable. It is double periodic with unit periods and can be represented by the double Fourier series

$$F(y, \xi) = \sum_{\lambda=(p,\theta) \in \Lambda} c_\lambda \exp(2\pi i(py + \theta\xi)),$$

where $\sum |c_\lambda|^2 = \|F\|^2$. By Proposition 3.1 the series $\sum c_\lambda \mathbf{e}_\lambda$ converges in $L_2(\mathbb{R})$ to a function g . On the other hand, because of (12),

$$Zf_\sharp(y, \xi) = Z\mathbf{e}_0 \sum c_\lambda \exp(2\pi i(py + \theta\xi)) = \sum c_\lambda Z\mathbf{e}_\lambda(y, \xi)$$

This implies $f_\sharp = g$, since Z is a unitary operator. Set $\gamma^\lambda(f) \doteq c_\lambda$ for $\lambda \in \Lambda$. ►

Lemma 6.3 *We have $F \in L_2(\mathbb{Q})$.*

PROOF OF LEMMA. By the previous Lemma the function $Zf_{\sharp}$ is ε -Hölder continuous in the interior of $\mathbb{Q}$. It vanishes at the sharp point, since $\gamma^{\sharp}(f_{\sharp}) = 0$. Therefore $|Zf_{\sharp}(z)| \leq C_{\delta} |z - 1/2 - \iota/2|^{\varepsilon}$. This implies

$$|F(z)| = \left| \frac{\exp(\pi y^2) Zf_{\sharp}}{\Theta(z)} \right| \leq C |z - 1/2 - \iota/2|^{\varepsilon-1},$$

since $\Theta(z)$ has simple zero at each sharp point. This implies the statement. ►

Remark 1. More strong inequality holds for $f \in H^{\delta}, \delta > 1$:

$$\sum_{\Lambda} (|\lambda| + 1)^{2\varepsilon} |\gamma^{\lambda}(f)|^2 \leq C_{\delta, \varepsilon} \|f\|_{\delta}^2, \quad 0 < \varepsilon < \delta - 1$$

Remark 2. We can take an arbitrary sharp point $\mu \in \Lambda + \sharp$ instead of $(1/2, 1/2)$ in Theorem 6.1.

7 Gabor coefficients

The coefficients are uniquely defined since of

Proposition 7.1 *If $\{c_{\lambda}\} \in l_2(\Lambda)$, $b \in \mathbb{C}$ and*

$$b\mathbf{e}_{\sharp} + \sum_{\Lambda} c_{\lambda} \mathbf{e}_{\lambda} = 0, \quad (20)$$

then $c_{\lambda} = 0$ for all λ .

PROOF. By Proposition 3.1 the series (20) converges in L_2 . Apply the Zak transform. By (12) we get

$$-bZ\mathbf{e}_{\sharp} = \sum c_{\lambda} Z\mathbf{e}_{\lambda} = Z\mathbf{e}_0 \sum c_{\lambda} \exp 2\pi\iota(p\xi + \theta y)$$

The series in the right-hand side converges to a function $g \in L_2(\mathbb{Q})$ since Z is a unitary operator. It follows $g = -bZ\mathbf{e}_{\sharp}/Z\mathbf{e}_0$. We have $Z\mathbf{e}_{\sharp}(1/2, 1/2) \neq 0$, whereas the function $Z\mathbf{e}_0(y, \xi) = \exp(-\pi y^2) \Theta(\xi + \iota y)$ vanishes in the point $1/2 + \iota/2$. Therefore the function g can not be square integrable unless $b = 0$. This yields $g = 0$, hence $c_{\lambda} = 0$ for all $\lambda \in \Lambda$. ►

Corollary 7.2 *We have the following explicit formula for the coefficients:*

$$\gamma_{\lambda}(f) = \int_0^1 \int_0^1 \exp(-2\pi\iota(py + \theta\xi)) \frac{\exp(\pi y^2) Zf_{\sharp}(y, \xi)}{\Theta(\xi + \iota y)} dy d\xi, \quad \lambda \in \Lambda \quad (21)$$

where $f_{\sharp} = f - \gamma^{\sharp}(f) \mathbf{e}_{\sharp}$.

Remark. It follows from a formula of M.Bastiaans [1], that the coefficients can be written in the form

$$\gamma^\lambda(f) = \langle f_\# | \exp(2\pi i \theta x) \gamma(x - p) \rangle, \quad \lambda = (p, \theta) \in \Lambda,$$

where

$$\gamma(x) = 2^{-1/4} (\pi/K_0)^{3/2} \exp(\pi x^2) \sum_{n+1/2 \geq x} (-1)^n \exp(-\pi(n+1/2)^2)$$

This function γ does not belong to L_2 , but the integrals $\langle f_\# | \exp(2\pi i \theta x) \gamma(x - p) \rangle$ converge since $f_\# \in H$ and $\gamma^\#(f_\#) = 0$.

8 Improving convergence

To ensure faster convergence of the relaxed Gabor series we impose more sharp conditions on f . Fix an integer $m > 0$ and consider the norm

$$\|f\|_{\delta, m} \doteq \left(\sum_{j=0}^m \|\mathbf{a}^j f\|_\delta^2 \right)^{1/2}$$

Let $H^{\delta, m}$ be the space of functions with finite norm $\|\cdot\|_{\delta, m}$.

Lemma 8.1 *For arbitrary different points $\mu_0, \dots, \mu_m \in \mathbb{C}$ the inverse to the VanderMond matrix $W = \{w_j^k \doteq (\mu_j)^k\}$ is equal to the matrix $V = \{v_j^k\}$ where*

$$\begin{aligned} v_j^k &\doteq \frac{\sigma_{m+1-j}^{(k)}}{p'(\mu_k)}, \quad p(\lambda) = \prod_{j=0}^m (\lambda - \mu_j), \\ \sigma_j^{(k)} &= (-1)^j \sigma_j(\mu_0, \dots, \widehat{\mu_k}, \dots, \mu_m) \end{aligned}$$

and σ_j denotes the j -th elementary symmetric polynomial.

PROOF is by straightforward check.

Lemma 8.2 *For an arbitrary natural m and any set of different points $\mu_0, \dots, \mu_m \in \Lambda + \#$ the are numbers $\{h_j^k\}$ such that*

$$\gamma^\#(\mathbf{a}^k \mathbf{d}_{m,j}) = \delta_j^k, \quad k, j = 0, \dots, m \quad (22)$$

where

$$\mathbf{d}_{m,j} = \sum_{s=0}^m h_j^s \mathbf{e}_{\mu_s}, \quad j = 0, \dots, m. \quad (23)$$

PROOF. Calculate the matrix

$$g_j^k \doteq \gamma^\sharp (\mathbf{a}^k \mathbf{e}_{\mu_j}), \quad j, k = 0, \dots, m$$

By (17) we have for an arbitrary $\mu = (q, \eta) \in \Lambda + \sharp$

$$\begin{aligned} Z\mathbf{e}_\mu(y, \xi) &= \exp \left(-\pi (y - 1/2)^2 + 2\pi \imath [y\eta + (q - 1/2)(\xi + 1/2)] \right) \\ &\quad \times \Theta(\xi + 1/2 + \imath(y - 1/2)) \end{aligned}$$

which yields $Z\mathbf{e}_\mu(\sharp) = \exp(\pi \imath \eta) \Theta(0)$. By (13)

$$\gamma^\sharp (\mathbf{a}^k \mathbf{e}_\mu) = \frac{\mu^k}{\imath \Theta(0)} Z\mathbf{e}_\mu(\sharp) = (-1)^{[\eta]} \mu^k$$

This yields $g_j^k = (-1)^{[\eta_j]} \mu_j^k$. The matrix $\{g_j^k\}$ is invertible and by Lemma 8.1 the entries of the inverse matrix are

$$h_j^k = (-1)^{[\eta_k]} \frac{\sigma_{m+1-j}^{(k)}}{p'(\mu_k)}, \quad k, j = 0, \dots, m.$$

Then (23) implies (22). $\blacktriangleright$

Lemma 8.3 *We have for any $\lambda \in \Phi$*

$$\left\| \sum \mathbf{d}_{m,j} \lambda^j \right\|_{\delta, m} \leq (M + 1)^{m+\delta} L^m \quad (24)$$

where

$$M \doteq \max |\mu_s|, \quad L = \max |\lambda - \mu_s|$$

PROOF. We have for $k, j = 1, \dots, m$

$$\left\| \mathbf{a}^k \sum_j \mathbf{d}_{m,j} \lambda^j \right\| = \left\| \sum_{j,s} h_j^s \lambda^j \mu_s^k \mathbf{e}_{\mu_s} \right\| \leq M^k \left| \sum_j h_j^s \lambda^j \right|$$

since $\|\mathbf{e}_\mu\| = 1$. Further

$$\sum_j h_j^s \lambda^j = \pm \sum_j \frac{\sigma_{m+1-j}^{(s)}}{p'(\mu_s)} \lambda^j = \pm \frac{p_s(\lambda)}{p_s(\mu_s)}$$

where $p_s(\lambda) \doteq \prod_{j \neq s} (\lambda - \mu_j)$. This yields

$$\left| \sum_j h_j^s \lambda^j \right| = \left| \frac{p_s(\lambda)}{p_s(\mu_s)} \right| = \prod_{j \neq s} \left| \frac{\lambda - \mu_j}{\mu_s - \mu_j} \right| \leq L^m$$

and (24) follows. $\blacktriangleright$

Theorem 8.4 *For an arbitrary natural m and different points $\mu_0, \dots, \mu_m \in \Lambda + \sharp$ there exists a family of continuous functionals γ_m^λ , $\lambda \in \Lambda$ in $H^m = \cup_{\delta > 1} H^{\delta, m}$ such that for an arbitrary $f \in H^m$ the equation holds in the space $W_2^k(\mathbb{R})$:*

$$f = \sum_{j=0}^m \gamma^\sharp(\mathbf{a}^j f) \mathbf{d}_{m,j} + \sum_{\Lambda} \gamma_m^\lambda(f) \mathbf{e}_\lambda, \quad (25)$$

where $\mathbf{d}_{m,j}$ are as in (23) and there exists for any $\delta > 1$ a constant C_δ such that

$$\sum_{\Lambda} (|\lambda|^2 + 1)^m |\gamma_m^\lambda(f)|^2 \leq C_\delta \|f\|_{\delta, m}^2. \quad (26)$$

PROOF. Set

$$f_\sharp = f - \sum_{j=0}^m \gamma^\sharp(\mathbf{a}^j f) \mathbf{d}_j \in H^m$$

By (14) and Lemma 6.2 the function $A^j Z f_\sharp$ is continuous in the interior of $\mathbb{Q}$ for $j \leq m$. We have $A = \bar{\partial}/2\pi i + y$, where $\bar{\partial} = (\partial_\xi + i\partial_y)$ and

$$A^j Z f_\sharp(y, \xi) = \Theta(\xi + iy) A^j [\exp(-\pi y^2) F(y, \xi)] = \Theta(\xi + iy) \exp(-\pi y^2) \bar{\partial}^j F(y, \xi),$$

since the function Θ is holomorphic and $A \exp(-\pi y^2) = 0$. Therefore the function $\bar{\partial}^j F$ is Hölder continuous in $\mathbb{Q}$ for $j \leq m$. The property (22) and (14) imply that $\gamma^\sharp(A^j Z f_\sharp) = 0$, $j = 0, \dots, m$ that is the left-hand side vanishes at the sharp point. By Lemma 6.3 the function

$$\bar{\partial}^j F = \frac{\exp(\pi y^2) A^j Z f_\sharp(y, \xi)}{\Theta(\xi + iy)}$$

is double periodic and belongs to $L_2(\mathbb{Q})$ for $j = 0, \dots, m$. Therefore $F \in W_2^m(\mathbb{R}^2/\Lambda)$ and the Fourier coefficients c_λ of F satisfy

$$\sum_{\Lambda} (|\lambda|^2 + 1)^m |c_\lambda|^2 \leq C \|F\|_{W_2^m}^2 \leq C'' \|f_\sharp\|_{\delta, m}^2 \leq C''' \|f\|_{\delta, m}^2$$

This yields (26) for $\gamma_m^\lambda(f) \doteq c_\lambda$, $\lambda \in \Lambda$. ►

A representation like (25) is unique, in spite of the additional terms:

Proposition 8.5 *If for some integer $m \geq 0$ the series*

$$\sum_{\Lambda} (|\lambda|^2 + 1)^m |c_\lambda|^2 \quad (27)$$

converges and

$$\sum_{j=0}^m b_j \mathbf{d}_j + \sum_{\Lambda} c_\lambda \mathbf{e}_\lambda = 0 \quad (28)$$

for some $b_0, \dots, b_m$, then $b_0 = \dots = b_m = 0$ and $c_\lambda = 0$ for all λ .

PROOF. The series in (28) converges to a function $g \in W_2^m(\mathbb{R})$. Moreover g belongs to the domain of the operators $\mathbf{a}^j, j = 1, \dots, m$. This follows from convergence of (27) and arguments of Proposition 3.1. By (14) the Zak transform Zf belongs to the domain of operators $A^j, j = 1, \dots, m$. By (12) we have

$$-\sum_0^m b_j Z\mathbf{d}_j = \sum_{\Lambda} c_{\lambda} Z\mathbf{e}_{\lambda} = Z\mathbf{e}_0 \sum_{\Lambda} c_{\lambda} \exp 2\pi i (p\xi + \theta y)$$

The series in the right-hand side converges to a function $g \in L_2(\mathbb{Q})$ and $g = -\sum b_j Z\mathbf{d}_j / Z\mathbf{e}_0$. We have $Z\mathbf{d}_0(z_{\sharp}) = 1$, whereas the functions $Z\mathbf{d}_1, \dots, Z\mathbf{d}_m$ and $Z\mathbf{e}_0(y, \xi)$ vanish at the sharp point. Moreover, we have $|Z\mathbf{e}_0(y, \xi)| \leq c|z - 1/2 - i/2|$. Therefore the function g can not be square integrable unless $b_0 = 0$. Similarly, the inclusion $Ag \in L_2(\mathbb{Q})$ implies that $b_1 = 0$ and so on. Therefore $b_0 = \dots = b_m = 0$ and $g = 0$; which yields $c_{\lambda} = 0$ for all $\lambda \in \Lambda$. ►

9 Gabor transform of a Gabor series

Let G be a set in the phase plane Φ and $r > 0$; for any $r > 0$ we denote by $G(r)$ the r -neighborhood of G . Consider a convergent series

$$g = \sum_{\lambda \in G} c^{\lambda} \mathbf{e}_{\lambda}$$

and estimate the Gabor transform of g :

Proposition 9.1 *For any subset $G \subset \Lambda$ and any $r > 0$ we have*

$$\int_{\Phi \setminus G(r)} |\langle g | \mathbf{e}_{\mu} \rangle|^2 d\mu \leq \exp(-\pi r^2) \sum |c^{\lambda}|^2$$

PROOF. By (4) we have for $\mu \in \Phi \setminus G(r)$

$$|\langle g | \mathbf{e}_{\mu} \rangle|^2 \leq \left(\sum_{\lambda \in G} |\langle \mathbf{e}_{\mu} | \mathbf{e}_{\lambda} \rangle| |c^{\lambda}| \right)^2 \leq \sum_G \exp(-\pi |\lambda - \mu|^2) \sum_G |c^{\lambda}|^2.$$

Integrating the right-hand side yields

$$\int_{\Phi \setminus G(r)} |\langle g | \mathbf{e}_{\mu} \rangle|^2 d\mu \leq \sum_{\lambda \in G} |c^{\lambda}|^2 \int_{\Phi} \exp(-\pi |\lambda - \mu|^2) \phi(\mu) d\mu.$$

where ϕ is the indicator function of the set $\Phi \setminus G(r)$. Write the sum over λ as the integral over Φ with the singular measure $\sigma(\lambda) = \sum_G |c^{\lambda}|^2 \delta_{\lambda}$. and change the

variable λ by $\kappa = \lambda - \mu$ in the double integral

$$\begin{aligned} \sum_{\lambda \in G} \int_{\Phi} \exp(-\pi |\lambda - \mu|^2) h(\mu) d\mu |c^\lambda|^2 &= \int_{\Phi} \int_{\Phi} \exp(-\pi |\lambda - \mu|^2) \phi(\mu) d\mu \sigma(\lambda) \\ &= \int_{\Phi} \exp(-\pi |\kappa|^2) \int_{\Phi} \phi(\mu) \sigma(\kappa + \mu) d\mu \end{aligned}$$

The interior integral in the right-hand side vanishes if $|\kappa| < r$, otherwise it is bounded by the total integral

$$\int \sigma(\kappa + \mu) d\mu = \sum |c^\lambda|^2 d\kappa$$

This yields the estimate for the right-hand side

$$\sum |c^\lambda|^2 \int_{|\kappa| \geq r} \exp(-\pi |\kappa|^2) d\kappa = \sum |c^\lambda|^2 \exp(-\pi r^2). \blacktriangleright$$

10 A ‘certainty’ theorem

Theorem 10.1 *Let $K \subset D$ be arbitrary bounded domains in Φ , such that D is r -neighborhood of K for some $r \geq r_0$. An arbitrary function $f \in H$ can be written in the form*

$$f = \sum_{\lambda \in \Lambda \cap D} \alpha^\lambda \mathbf{e}_\lambda + \sum_{\mu \in (\Lambda + \sharp) \cap D \setminus K} \omega^\mu \mathbf{e}_\mu + \phi_r$$

where for $f \in H^\delta$

$$\|\phi_r\| \leq \left(\int_{\Phi \setminus D} |\langle f | \mathbf{e}_\mu \rangle|^2 d\mu \right)^{1/2} + C_\delta r^\delta \exp(-r/e) \|f\|_\delta. \quad (29)$$

and the constant C_δ does not depend on K and D .

Remark 1. Note that

$$\alpha^\lambda = \gamma^\lambda(f) + O(r^\delta \exp(-r/e))$$

for $\lambda \in K$. The geometry of the two first terms is shown in Fig.1.




Fig.1: The lattice Λ , domains K (thin lines) and D (thick lines) in the phase plane.

Remark 2. The exponential term in the estimate (29) is indispensable, but its form might be made sharper. Indeed, vanishing of Gabor means $\langle f | \mathbf{e}_\mu \rangle$ for $\mu \in \Phi \setminus D$ does not guarantee that f can be represented by a Gabor functions supported in $\Lambda \cap D$ without additional term. This follows from Proposition 9.1 where the exponential factor $\exp(-\pi r^2)$ is sharp.

PROOF OF THEOREM. Set $K_+ = K(l)$, $U = K(r/2)$, $D_- = K(r-l)$ where the parameter $l = O(r^{1/2})$ will be specified later. We have $K \subset K_+ \subset U \subset D_- \subset D$. By Theorem 6.1 we can write $f = f_U + g$, where

$$f_U \doteq \sum_{\Lambda^\# \cap U} \gamma^\lambda(f) \mathbf{e}_\lambda, \quad g \doteq \sum_{\Lambda \setminus U} \gamma^\lambda(f) \mathbf{e}_\lambda. \quad (30)$$

and the sharp point $\sharp$ is chosen in $U \setminus K$. Estimate Gabor transform of g . By Proposition 9.1 and Theorem 6.1

$$\int_{K_+} |\langle g | \mathbf{e}_\mu \rangle|^2 d\mu \leq \exp(-\pi(r/2 - l)^2) \sum_{\lambda \in \Lambda \setminus U} |\gamma^\lambda(f)|^2 \leq C \exp(-\pi(r/2 - l)^2) \|f\|_\delta^2$$

If $\mu \in \Phi \setminus D_-$, we can write $\langle g | \mathbf{e}_\mu \rangle = \langle f | \mathbf{e}_\mu \rangle - \langle f_U | \mathbf{e}_\mu \rangle$ since of (30). Apply again Proposition 9.1 and Theorem 6.1 and obtain

$$\int_{\Phi \setminus D_-} |\langle g | \mathbf{e}_\mu \rangle|^2 d\mu \leq \int_{\Phi \setminus U_+} |\langle f | \mathbf{e}_\mu \rangle|^2 d\mu + \exp(-\pi(r/2 - l)^2) \|f\|_\delta^2,$$

since f_U is a sum of functions $\mathbf{e}_\lambda, \lambda \in U$. By Proposition 2.1 we can write

$$\begin{aligned} g &= \int_{K_+} \langle g | \mathbf{e}_\mu \rangle \mathbf{e}_\mu d\mu + \int_{\Phi \setminus D_-} \langle g | \mathbf{e}_\mu \rangle \mathbf{e}_\mu d\mu + \int_{D_- \setminus K_+} \langle g | \mathbf{e}_\mu \rangle \mathbf{e}_\mu d\mu \\ &\doteq g_+ + g_- + g_0 \end{aligned} \quad (31)$$

The norms of the first two terms are bounded as follows

$$\|g_+\|^2 \leq \exp(-\pi(r/2 - l)^2) \|f\|_\delta^2, \quad (32)$$

$$\|g_-\|^2 \leq \exp(-\pi(r/2 - l)^2) \|f\|_\delta^2 + \int_{\Phi \setminus D} |\langle f | \mathbf{e}_\mu \rangle|^2 d\mu \quad (33)$$

We fix a natural $m \leq r - 1$ and transform the integral g_0 as follows: for a point $\mu \in D_- \setminus K_+$ we choose a point $\lambda \in \Lambda \cap D \setminus K$ and such that $|\lambda - \mu| \leq 1/\sqrt{2}$ and consider the closed square $Q(\mu)$ centered at λ with side $\sqrt{m+1}$, see Fig.2.




Fig.2: λ is the closest point to μ in the lattice Λ (thick points) . Points of the set $(\Lambda + \sharp) \cap Q(\mu)$ are shown by $\sharp$;

The set $Q(\mu)$ contains $m+1$ different points $\mu_0, \dots, \mu_m \in \Lambda + \sharp$ and is contained in the l -neighborhood of μ where $l = ((m+1)/2)^{1/2} + 1$. It follows that $Q(\mu) \subset D \setminus K$ and

$$\max |\mu - \mu_s| \leq l, \max |\lambda - \mu_s| \leq l. \quad (34)$$

Applying Theorem 8.4 to the function $\phi = \mathbf{e}_{\mu-\lambda}$ and sharp points $\mu_0 - \lambda, \dots, \mu_m - \lambda$ yields

$$\begin{aligned} \mathbf{e}_{\mu-\lambda}(x) &= \sum_{j=0}^m \gamma^\sharp(\mathbf{a}^j \mathbf{e}_{\mu-\lambda}) \mathbf{d}_{m,j} + g_{\mu,\lambda}, \\ g_{\mu,\lambda} &\doteq \sum_{\Lambda} \gamma_m^\nu(\mathbf{e}_{\mu-\lambda}) \mathbf{e}_\nu, \end{aligned} \quad (35)$$

where $\mathbf{d}_j$ belong to the linear span of $\mathbf{e}_{\mu_s-\lambda}, j = 0, \dots, m$. We have

$$\|g_{\mu,\lambda}\|_{\delta,m} \leq \|\mathbf{e}_{\mu-\lambda}\|_{\delta,m} + \left\| \sum \gamma^\sharp(\mathbf{a}^k \mathbf{e}_{\mu-\lambda}) \mathbf{d}_{m,k} \right\|_{\delta,m}.$$

and $\mathbf{a}^k \mathbf{e}_{\mu-\lambda} = (\mu - \lambda)^k \mathbf{e}_{\mu-\lambda}$. By Proposition 8.3 and (34)

$$\begin{aligned} \left\| \sum_k \gamma^\sharp(\mathbf{a}^k \mathbf{e}_{\mu-\lambda}) \mathbf{d}_{m,k} \right\|_{\delta,m} &\leq C(l+1)^{m+\delta} l^m \leq C(m+1)^{m+\delta} \\ \|\mathbf{e}_{\mu-\lambda}\|_{\delta,m} &\leq C(l+1)^{m+2} \leq C'(m+1)^{(m+\delta)/2} \end{aligned}$$

where the constants does not depend on m . This yields

$$\begin{aligned} \left\| \sum \gamma^\sharp(\mathbf{a}^k \mathbf{e}_{\mu-\lambda}) \mathbf{d}_{m,k} \right\|_{\delta,m} &\leq C(m+1)^{m+\delta}, \\ \|g_{\mu,\lambda}\|_{\delta,m} &\leq C'(m+1)^{m+\delta} \end{aligned}$$

By Theorem 8.4

$$\sum (|\nu|^2 + 1)^m |\gamma_m^\nu(\mathbf{e}_{\mu-\lambda})|^2 \leq C(m+1)^{2(m+\delta)} \|\mathbf{e}_{\mu-\lambda}\|_{\delta,m}^2, \quad (36)$$

Let $\lambda = (p, \theta) \in \Lambda$; apply the operator T_λ as in (9). For $\mu = (q, \xi), \nu = (r, \eta)$ we obtain

$$T_\lambda \mathbf{e}_{\mu-\lambda} = \exp(-2\pi i p(\xi - \theta)) \mathbf{e}_\mu, \quad T_\lambda \mathbf{e}_\nu = \exp(-2\pi i p\eta) \mathbf{e}_{\nu+\lambda}$$

and

$$\begin{aligned} \mathbf{e}_\mu(x) &= \sum_{j=0}^m \gamma^\sharp(\mathbf{a}^j \mathbf{e}_{\mu-\lambda}) \exp(2\pi i p(\xi - \theta)) \mathbf{d}_{j,m} \\ &\quad + \sum_{\Lambda} \gamma_m^\nu(\mathbf{e}_{\mu-\lambda}) \exp(2\pi i p(\xi - \theta + \eta)) \mathbf{e}_{\nu+\lambda} \end{aligned} \quad (37)$$

The function $\mathbf{d}_{j,m}$ belongs to the linear envelope of $\mathbf{e}_\kappa, \kappa \in \Lambda + \sharp$. Change the variable $\nu + \lambda$ by ν in the second sum and write this equation in the form

$$\mathbf{e}_\mu(x) = \sum_{\kappa \in (\Lambda + \sharp) \cap Q(\mu)} \beta_\mu^\kappa \mathbf{e}_\kappa + \sum_{\lambda \in \Lambda \cap D} \varepsilon_\mu^\lambda \mathbf{e}_\lambda + \sum_{\nu \in \Lambda \setminus D} \omega_\mu^\nu \mathbf{e}_\nu \quad (38)$$

where $\omega_\mu^\nu = \gamma_m^{\nu-\lambda}(\mathbf{e}_{\mu-\lambda}) \exp(2\pi i r(\xi - \theta + \eta))$. Estimate the third term by means of (36)

$$\begin{aligned} \sum_\nu |\omega_\mu^\nu|^2 &= \sum_{\lambda \in \Lambda \setminus D} |\gamma_m^{\nu-\lambda}(\mathbf{e}_{\mu-\lambda}) \exp(2\pi i r(\xi - \theta + \eta))|^2 \\ &\leq C(m+1)^{2m+2} (|\nu - \lambda| + 1)^{-2m} \leq C(m+1)^{2(m+\delta)} r^{-2m} \end{aligned}$$

since $|\nu - \lambda| \geq r-2^{1/2}$. Taking $m = r/e - 1$ we obtain $(m+1)^{m+\delta} r^{-m} \leq C r^\delta \exp(-r/e)$. The relation $l = O(m^{1/2}) = O(r^{1/2})$ is fulfilled for this choice of m . Taking in account that the square $Q(\mu)$ is contained in $D \setminus K$ and integrating (38) on $D_- \setminus K_+$ against the density $\langle g | \mathbf{e}_\mu \rangle d\mu$, yields

$$g_0 = \sum_{\kappa \in (\Lambda + \sharp) \cap D \setminus K} \beta^\kappa \mathbf{e}_\kappa + \sum_{\lambda \in \Lambda \cap D} \varepsilon^\lambda \mathbf{e}_\lambda + \sum_{\lambda \in \Lambda \setminus D} \omega^\nu \mathbf{e}_\nu,$$

where

$$\beta^\kappa = \int_{D_- \setminus K_+} \beta_\mu^\kappa \langle g | \mathbf{e}_\mu \rangle d\mu, \quad \varepsilon^\lambda = \dots, \quad \omega^\nu = \dots$$

By Proposition 3.1 and Theorem 6.1 we have

$$\left(\sum |\omega^\nu|^2 \right)^{1/2} \leq C \exp(-r/e) \left(\int |\langle g | \mathbf{e}_\mu \rangle|^2 d\mu \right)^{1/2} \leq C r^\delta \exp(-r/e) \|g\| \quad (39)$$

$$\leq C_\delta r^\delta \exp(-r/e) \|f\|_\delta \quad (40)$$

Finally we get

$$f = \sum_{\lambda \in \Lambda \cap D} \alpha^\lambda \mathbf{e}_\lambda + g_+ + g_- + \sum_{\nu \in (\Lambda + \sharp) \cap D \setminus K} \beta^\nu \mathbf{e}_\nu + \sum_{\lambda \in \Lambda \setminus D} \omega^\nu \mathbf{e}_\nu$$

where $\alpha^\lambda = \gamma^\lambda(f) + \varepsilon^\lambda$ for $\lambda \in \Lambda \cap U$ and $\alpha^\lambda = \varepsilon^\lambda$ for $\lambda \in \Lambda \cap D \setminus U$ and the term $\gamma^\sharp(f) \mathbf{e}_\sharp$ is included in the second sum. We arrange this sum as follows

$$f = \sum_{\Lambda \cap D} \alpha^\lambda \mathbf{e}_\lambda + \sum_{\kappa \in (\Lambda + \sharp) \cap D \setminus K} \beta^\kappa \mathbf{e}_\kappa + \phi_r$$

where and

$$\phi_r = g_+ + g_- + \sum_{\lambda \in \Lambda \setminus D} \omega^\lambda \mathbf{e}_\lambda. \quad \blacktriangleright$$

By (32), (33) and (40) we have

$$\begin{aligned}\|\phi_r\| &\leq \|g_+\| + \|g_-\| + \left\| \sum \omega^\nu \mathbf{e}_\nu \right\| \\ &\leq \left(\int_{\Phi \setminus D} |\langle f | \mathbf{e}_\mu \rangle|^2 d\mu \right)^{1/2} + 2 \exp(-\pi(r/2 - l)^2/2) \|f\|_\delta + C_\delta r^\delta \exp(-r/e) \|f\|_\delta,\end{aligned}$$

which yields (29) for any sufficiently large r . ►

11 Metaplectic group

Remind that the Weyl-Heisenberg group is the space $X \times X^* \times \mathbb{R}$ with the group operation

$$(x, \xi, \tau) \cdot (x', \xi', \tau') = \left(x + x', \xi + \xi', \tau + \tau' + \frac{1}{2}(x'\xi - x\xi') \right).$$

Definition. A linear transform S of Φ is called *simplectic*, if it preserves the canonical bilinear form $\sigma[(x, \xi), (y, \eta)] = \eta x - \xi y$. We can see that $\sigma[u, v] = \langle u | Jv \rangle$, where

$$J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

In particular, $\lambda = (p, \theta) \mapsto J\lambda = (\theta, -p)$ is a linear simplectic transform. In the case $X = \mathbb{R}$ a linear transformation S in $\mathbb{R} \times \mathbb{R}^*$ is simplectic if the matrix (a, b, c, d) of S satisfies $ad - bc = 1$. For an arbitrary linear simplectic transformation S such that $b \neq 0$ the integral transform

$$M_S f(x) = (\imath b)^{-1/2} \int \exp \left(\pi \imath \left(\frac{d}{b} x^2 - \frac{2}{b} yx + \frac{a}{b} y^2 \right) \right) f(y) dy.$$

is well defined operator in L_2 . If $n = 0$, the operator M_S is defined as composition $M_T F$, where F is the Fourier transform and $T = \imath^{1/2} S J^{-1}$. The operator M_S has unitary closure in L_2 . The equation $M_S M_T = \pm M_{ST}$ holds for any simplectic transformations S, T . It is called metaplectic (two-valued) representation of S . In particular, the metaplectic operator M_J is the Fourier transform F times the factor $\imath^{1/2}$. This representation is single-valued on the irreducible two-fold covering of the group S . See more information in [5].

Any rotation $S(x, \xi) = (\cos \varphi x - \sin \varphi \xi, \sin \varphi x + \cos \varphi \xi)$ is an orthogonal simplectic transformation.

Proposition 11.1 *If S is a rotation as above, then*

$$\begin{aligned}M_S \mathbf{a} &= \exp(-\imath \varphi) \mathbf{a} M_S \\ M_S \mathbf{a}^+ &= \exp(\imath \varphi) \mathbf{a}^+ M_S\end{aligned}$$

where $\mathbf{a}^+, \mathbf{a}$ is the creation and the annihilation operator, respectively.

PROOF. Direct calculation.

Proposition 11.2 *For any rotation S in the phase plane and any point λ we have*

$$M_S(\mathbf{e}_\lambda) = \pm \exp(\imath\varphi/2) \exp(\pi\imath(p\theta - q\eta)) \mathbf{e}_{S(\lambda)} \quad (41)$$

where $(q, \eta) = S(p, \theta)$

PROOF is straightforward.

Remark. The equation (41) means that M_S transforms a Gabor function $\mathbf{e}_\lambda$ to another Gabor function (up to a phase factor) while the ‘quantum support’ λ of a Gabor function moves by action of the corresponding geometric transform S . In particular, the Fourier transform belongs to the metaplectic group: $F = M_J$. We have for any $\lambda = (p, \theta)$, $\hat{\mathbf{e}}_\lambda(\eta) = \pm \exp(\imath\varphi/2) \mathbf{e}_{\hat{\lambda}}(\eta)$, $\hat{\lambda} = J\lambda = (\theta, -p)$ for a real ϕ . The metaplectic representation can be thought as a ‘quantization’ of group of symplectic transforms.

Corollary 11.3 *For any $\delta > 1$ and natural m the space $H^{\delta,m}$ is invariant under action of the metaplectic representation of the rotation group.*

PROOF. For any rotation S we have

$$\langle f | \mathbf{e}_\lambda \rangle = \langle M_S f | M_S \mathbf{e}_\lambda \rangle = \exp(\imath\varphi/2) \langle M_S f | \mathbf{e}_\lambda \rangle$$

which yields $|\langle M_S f | \mathbf{e}_\lambda \rangle| = |\langle f | \mathbf{e}_\lambda \rangle|$. It follows that H^δ is invariant. The same true for $H^{\delta,m}$ since of (41). ►

Corollary 11.4 *For an arbitrary orthogonal symplectic transformation S in the phase space, theorem 6.1 holds for functions $f \in H$ and Gabor system $\mathbf{e}_\lambda, \lambda \in S(\Lambda^\sharp)$.*

Remark. The operator M relates to the metaplectic representation μ in the sense of [5] by the equation $M_S = \mu(\tilde{S})$, where the symplectic matrix $\tilde{S}$ is obtained from S by changing sign at b and c and replacing θ by $-\theta$. This results in the representation μ which is chosen in [5]. Note that the mapping $S \mapsto \tilde{S}$ is a involution in the symplectic group.

References

- [1] M.J. BASTIAANS, A sampling theorem for the complex spectrogram, and Gabor expansion of a signal in Gaussian elementary signals. Optical Engineering 20:4 (1981), 594-598.

- [2] I. DAUBECHIES, Ten lectures on wavelets. Philadelphia, 1992.
- [3] H.G. FEICHTINGER, K. GRÖCHENIG, Gabor frames and time-frequency analysis of distributions, J. of Functional Analysis 146 (1997), 464-495.
- [4] H.G. FEICHTINGER, T. STROHMER (eds), Gabor analysis and algorithms. Theory and Applications, Applied and Numerical Harmonic Analysis, Birkhäuser, 1998.
- [5] G.B. FOLLAND, Harmonic analysis in phase space. Princeton Univ.Press, Princeton, New Jersey, 1989.
- [6] D. GABOR, Theory of communication. J. Inst. of El. Eng. 93:3 (1946), 429-457.
- [7] A.J.E.M. JANSSEN, Gabor representation of generalized functions. J. of Math. Anal. and Appl. 83 (1984), 377-394.
- [8] H.J. LANDAU, H.O. POLLAK, Prolate spheroidal wave functions, Fourier analysis and uncertainty, II; III. Bell System Tech.J. 40 (1961), 65-84; 41 (1962), 1295-1336.
- [9] YU.I. LYUBARSKII, Frames on the Bargmann space of entire functions, Adv. Soviet Math. 11 AMS, Providence RI, 1992, pp.167-180.
- [10] YU.I. LYUBARSKII, K. SEIP, Convergence and summability of Gabor expansion at the Nyquist density, Journal of Fourier Anal. and Appl. 5:2/3 (1999), 127-157.
- [11] V.I. SMIRNOV, A course of higher mathematics. V, 1964, Pergamon.